

FOR VETERINARY USE ONLY



KEEP OUT OF REACH
OF CHILDREN

PRECAUTIONS

Although Piperazine is a drug with a wide margin of safety, occasionally an animal may show nausea, vomiting, or muscular tremors. Such side effects are usually associated with over-dosage. Therefore, recommended dosage should be followed carefully. Animals with known kidney pathology should be treated only by a veterinarian.

FOR ORAL ANIMAL USE ONLY
NOT FOR HUMAN CONSUMPTION

KEEP OUT OF REACH OF CHILDREN

STORE OUT OF DIRECT SUNLIGHT AND
ABOVE FREEZING TEMPERATURES.

Consult your veterinarian for assistance in the
diagnosis, treatment and control of parasitism.

NDC 30798-254-31

Piperazine-17 Medicated

A wormer for use in drinking water for the control of large roundworms (*Toxocara canis* and *Toxascaris leonina*) in dogs and cats, and large roundworms (*Parascaris equorum*), strongyles (*Strongylus vulgaris*) and small strongyles and pinworms (*Oxyuris equi*) in horses.

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE ADDITIONAL PRECAUTIONS ON SIDE PANELS

Each 100 cc contains 17 grams of Piperazine base. Color, flavoring, and preservative added.

NET CONTENTS: 16 fl. oz. (1 PT.) (473 mL)

DIRECTIONS FOR USE

DOGS AND CATS - Give 1 ml. (15 drops) PIPERAZINE - 17 for each 6 pounds of body weight. Dose the dog and cat individually or give in milk, water or food. Repeat treatment in 30 days or when necessary.

HORSES: (non-food producing) Give 1 fluid ounce PIPERAZINE - 17 for each 100 pounds of bodyweight. Give the full dose with a dose syringe. If feed administration is preferred, give bran mash with a small amount of PIPERAZINE - 17 for 2 or 3 feedings, then give full dose in bran mash. Repeat treatment in 30 days or when necessary.

1-1561

Rev. 11-00

Manufactured for:
DURVET, INC.
Blue Springs, Missouri 64014

Lot No.

Exp. Date

