Safe Guard® Paste 10%

Equine Dewormer

INDICATIONS:
Safe-Guard® (fenbendazole) Paste 10% is indicated for the control of large strongyles *(Strongylus edentatus, S. equinus, S. vulgaris)*, encysted early third stage (hypobiotic), late third stage and fourth stage cyathostome larvae, small strongyles, pinworms *(Oxyuris equi)*, ascarids *(Parascaris equorum)*, and arteritis caused by fourth stage larvae of *Strongylus vulgaris* in horses.
Safe-Guard® (fenbendezole) Paste 10% is approved for use concomitantly with an approved form of trichlorfon. Trichlorfon is approved for the treatment of stomach bots *(Gasterophilus* spp.) in horses. Refer to the manufacturer's label for directions for use and cautions for trichlorfon.

FOR USE ON:
Horses

COMPOSITION:
fenbendazole…….100mg

DOSAGE and ADMINISTRATION:
Safe-Guard® (fenbendazole) Paste 10% is administered orally at a rate of 2.3 mg/lb (5mg/kg) for the control of large strongyles, small strongyles, and pinworms. For control of encysted early third stage (hypobiotic), late third stage and fourth stage cyathostome larvae, and fourth stage larvae of *Strongylus vulgaris,* the recommended dose is 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days.

DIRECTIONS FOR USE:
1. Determine the weight of the horse.
2. Remove syringe tip.
3. Turn the dial ring until the edge of the ring nearest the tip lines up with zero.
4. Depress plunger to advance paste to tip.
5. Now set the dial ring at the graduation nearest the weight of the horse (do not underdose).
6. Horse's mouth must be free of food.
7. Insert nozzle of syringe through the interdental space and deposit the paste on the back of the tongue by depressing the plunger.

PACKAGING:
25 gram paste, 36 tubes per case UPC # 0-21784-471-25 1

Distributed by: durvet

DURVET, INC. ANIMAL HEALTH PRODUCTS
P.O. Box 279 · Blue Springs, Missouri 64014 · 800/821-5570 · 816/229-9101
Fax 816/224-3080 · E-mail: info@durvet.com · www.durvet.com