

DURVAC LEPTO

5

Leptospira Canicola-
Grippotyphosa Hardjo-
Icterohaemorrhagiae-
Pomona Bacterin



Indications

For use in healthy cattle and swine as an aid in the prevention of leptospirosis caused by the micro-organisms named.

Contents

Product is aluminum hydroxide adsorbed.

Directions

Shake well before use. Use entire contents when first opened.

Dosage & Administration

Inject 2 ml intramuscularly or subcutaneously. For swine, a second dose should be administered 2-4 weeks later. Annual single dose re-vaccination is recommended for both species.

Precaution(s)

Store at 2° to 7° C. Do not freeze.

Caution(s)

Anaphylactoid reaction may occur following administration of products of this nature. If noted, administer adrenalin or equivalent.

Warning(s)

Do not vaccinate within 21 days before slaughter.
For use in animals only.
FOR VETERINARY USE ONLY.

Manufactured By

Colorado Serum Co.
4950 York St.
Denver, CO 80216
U.S. Vet. Lic. No. 188

Packaging

10 dose (20 mL) vials,
UPC# 7-45801-11852-2
50 dose (100 mL) vials,
UPC# 7-45801-11854-6

081017



Durvet, Inc. • P.O. Box 279, Blue Springs, MO 64014
(800)821-5570 • (816)229-9101 • Fax (816)224-3080
info@durvet.com • www.durvet.com