

# Pyrante S TYPE C MEDICATED FEED



## **SWINE ANTHELMINTIC**

#### Indications:

Pyrantel-S Type C Medicated Feed is indicated for removal and control of large roundworm (Ascaris suum) and nodular worm (Oesophagostomum) infections in swine.

#### Approved For Use In:

Swine.

#### **Product Advantages:**

- Easy to feed pellet form
- Scoop included measures approximately 0.50 pound (8 oz).
- Removes and controls large roundworms and nodular worms.

#### **Active Ingredient:**

Pyrantel Tartrate......0.0881% (800 grams per ton)

### Ingredients:

Grain products, processed grain by-products, plant protein products, calcium carbonate, sodium chloride, monocalcium phosphate, L-Lysine, roughage products, iron sulfate, zinc sulfate, copper sulfate, manganous oxide, dried Pichia pastoris fermentation extract, vitamin E supplement, niacin supplement, d-calcium pantothenate, sodium selenite, vitamin A acetate, riboflavin supplement, menadione nicotinamide bisulfite (source of vitamin K and niacin), vitamin B12 supplement, vitamin D<sub>3</sub> supplement and ethylenediamine dihydriodide.

#### Packaging:

5 lb bag, 10 per bailer bag, UPC# 7-45801-07971-7



**Contains Banminth®** (Pyrantel Tartrate)

#### **Directions for Use:**

- Enclosed scoop measures approximately 0.50 pound (8 oz).
- Feed as the sole ration for a single therapeutic treatment at the rate of 1 pound of feed per 40 pounds of body weight for animals up to 200 pounds. For animals 200 pounds or over, feed 5 pounds of feed per head. Pigs will generally consume this feed within a few hours, after which regular feeding should be resumed.
- Better worming is achieved when pigs are separated by size into different lots or pens for treatment.
- Water should be made available to animals during fasting and treatment periods.
- Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.



www.durvet.com

081817