

WARNING - NOT FOR USE IN HUMANS.
Keep this and all drugs out of the reach of children.

RESIDUE WARNING: Do not treat cattle within 35 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not treat swine within 18 days of slaughter.

PRECAUTIONS: Use automatic syringe equipment only. For subcutaneous injection in cattle and swine **only**. Ivermectin (ivermectin) Injection has been developed specifically for use in cattle, swine, reindeer, and American bison **only**. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

NDC 30798-844-05

durvet

IVERMECTIN
(ivermectin)
INJECTION for Cattle & Swine



1% Sterile Solution
Restricted Drug (California) -
Use Only as Directed.

ANADA 200-447, Approved by FDA
Net Contents: 50 mL
(1.69 fl oz)

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

INDICATIONS

Cattle: For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs (note insert precautions), sucking lice, and mange mites in cattle.

Swine: For the treatment and control of gastrointestinal roundworms, lungworms, lice, and mange mites in swine.

RECOMMENDED DOSE

Cattle: 1 mL per 110 lb bodyweight

Swine: 1 mL per 75 lb bodyweight

See package insert for complete indications, precautions, warnings, residue information, and use directions.

Do not contaminate water. Dispose of containers in an approved landfill or by incineration.

Store at 20°C to 25°C (68°F to 77°F).

Protect from light.

Manufactured for:
Durvet, Inc.
Blue Springs, MO 64014
www.durvet.com

Manufactured by:
Bimeda-MTC Animal Health Inc.
Cambridge, ON N3C 2W4
www.bimeda.com

TAKE TIME



OBSERVE LABEL
DIRECTIONS

8IVE045

Rev: 02/16