Dairy Bomb 55

Indications:

Dairy Bomb 55 is indicated for use in the control of flying and crawling insects Beef Cattle Operations, Dairy Barns (including Milk Houses and Milk Rooms), Hog Houses, Barns, Stables, Animal Quarters, Milk Rooms, and Poultry Houses. (see label for extensive list)

Approved For Use On:

Cattle, Horses and Swine.

Approved for use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food-Handling Plants, Dairies, Milk Houses, Cattle, Horse, Poultry and Swine Quarters.

Active Ingredients:

Pyrethrins	0.5%
Piperonyl Butoxide*	
OTHER INGREDIENTS**	94.5%
TOTAL1	

*(Butylcarbityl)(6-Propylpiperonyl) ether and related compounds **(Contains Petroleum Distillates: 1,1-Difluroethane)

Packaging:

25 oz. aerosol, 6 cans/case UPC# 7-45801-33303-1





DAIRY BOMB 55

040825



Net Contents: 25 oz. (709 grams)

KEEP OUT OF REACH OF CHILDREN CAUTION

"Betweet to WIDE-Propyloperory) either and related compounds
"Contains Petrideum Distillates; 1,1-Lhfluroethane)
EPA Flag, No. 47000-69-12281

OTHER INGREDIENTS**

TOTAL

ACTIVE INGREDIENTS: Piperanyl Butaxide*.

n nuts, dried fruit, bear s, dried prunes, dried

OR USE ON STORED PRODUCTS

Have product container or label with you when calling a poison control center, wy ay mouth to an unconscious person. Unless told to do so by the poison control off contaminated clothing skin innmediately with plenty of water for 15-20 minutes. poison control center or doctor for further treatment advi doctor, or going for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas). PRECAUTIONARY STATEMENTS HAZAROS TOHUMANG AND OMESTIC ANIMALS CAUTORY FIRING IN ASSAULT ANIMALS If on Skin or on Clothing:

Ann treatment to completed or conscions, warenchings, zowhangs, priefers, and contriportes: pray this product throughly into insect hinding places such as a product throughly into insect hinding places and ands, under intropractics, along baseboards, and in chrouge reas Spray directly on insects where possible Repeat as

r ants' trails, nests, and points of entry. Spray directly or where possible. Repeat no more than once per day.

bins and trucks, cargo

OR USE IN FOOD STORAGE SITES

en leave the treated areas. Keep 5 minutes. Ventilate the room

oom with mist, and n closed for at least

n skin. Avoid contact with skin, eyes or clothing. Prolonged or ause allergic reaction in some individuals.

Personal Protective Equipment (PPE)

ig pants, loes and socks, and nemical-resistant glover

s (such as barrier laminate, nitrile rubber, neoprene rubber, or Viton). er Salety Requirements

rorm other laundry. Discard clothin ontarrinated with this product's tons for clearing/maintaining PPE. If no such instructons for was vater Keep and wash PPE separately from other laundy. Dissard o s that have been drenched or heavily contarrinated with this produ

rash hands before eating, drinking, chewing guri, using tibaaco, or using the toilet errove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and j

s should remove PPE immediately after handling this product. Wash the outside of gloves b wing them. As soon as possible, wash throroughly and change into clean clothing.

YSICAL OR CHEMICAL HAZARDS: Combustble. Contents under pressure. Keep away from a sparks, and heated surfaces. Do not pureture or incinerate container. Exposure to temperature

varnished surfaces or directly into electronic equipment such as i

DIRECTIONS FOR USE

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions, and prec with its labeling. violation of Federa

f the debris should be removed and burned to kill eggs needs that might be present. Before processing begins

upright, nozzle directed away from you, and press actuator button and spray as directed **JSE RESTRICTIONS:**

restbek, do not enter or allow others to enter treated areas until vapors, mists, aread and the areas have been thoroughly ventilated.

cations, cover or remove food. cal facilities, remove patients prior to treatment. Ventilate treated areas for

v returning patients. Indeor food handling and processing facilities, do not make space spray applications.

s space spray applications, cover food processing surfaces or clean after treatment and before Do not apply more than once per day.



rus or facilities: Close mik bulk tank ation from spray and from dead or r cover miking utensils before appli-

Dairy Bomb-55

FIRST AID

A pest, birds leventin continencial poul by operational, ter fish aquanums before spraying, v ues only.

or use only. use in outdoor residential misting systems.

COR USE IN HOME AND FOOD PROCESSING AREAS For files, mosquirees, small flying monts (millers), Asian lady beetles ", and gnats:

DAIRY BOMB 55

040825