

FlyRID[®] PLUS

With
CITRONELLA

Description:

FlyRID[®] Plus is a multi-species, multi-purpose insect control spray that provides powerful yet safe protection against harmful pests. This ready-to-use permethrin product is for use directly on animals and their premises, as well as other indoor/outdoor applications.

For Use On:

Horses, dogs, goats, sheep, hogs, beef and dairy cattle and their premises, such as barns, stables, pet beds, porches, poultry premise sites and garages. Also approved for other indoor-outdoor applications as indicated by label directions.

Product Advantages:

- Ready-to-use spray that's safe, effective and economical
- Permethrin is a stable chemical used in insect control
- Contains citronella
- Can be used on Horses, Dogs, Goats, Sheep, Hogs, Beef and Dairy Cattle
 - Controls horse flies, horn flies, face flies, house flies, stable flies, deer flies, mosquitoes, and gnats
- Great for barns, stables and other premises
- Controls fleas and ticks on dogs for 35 days
 - Controls ticks, which may carry lyme disease (deer tick, lone star tick, brown dog tick, American dog tick and western black legged tick).
- Provides 60-day control of cockroaches, spiders, ants, crickets, scorpions, millipedes and centipedes.
- Controls bed bugs and mites in poultry houses.

Active Ingredients:

Permethrin.....0.5%



Packaging:

32 oz. spray bottle, 12 bottles per case
UPC# 7-45801-31034-6

Gallon bottle, 4 bottles per case
UPC# 7-45801-31036-0



072926

FIRST AID

IF ON SKIN OR ON CLOTHING:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for further treatment advice.
Have product contained or label with you when calling a poison control center, or doctor, or going in for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-423-0318 (all other areas).	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

- Appliers and other handlers must wear the following:
 - Long-sleeved shirt and long pants;
 - Shoes and socks; and
 - Chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or Viton.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees or other pollinating insects are foraging the treatment area.

Durvet®, the Durvet logo and FlyRID® are registered trademarks of Durvet, Inc. All rights reserved.

Manufactured in the U.S.A.
with global components

OPEN HERE

ISS26X005 EEA Reg. No. 47000-148-12281

EEA Est. No. 47000-14-01

7

4580131034

6

PERMETHRIN GROUP 3A INSECTICIDE



durvet

FlyRID[®] PLUS



Multi-Purpose Spray

Controls Flies, Mosquitoes and Gnats
35-Day Control of Fleas & Ticks on Dogs
With **CITRONELLA**

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See back panel for additional
Precautionary Statements

NET CONTENTS: 1 Qt. (946 mL)

ACTIVE INGREDIENT:

Permethrin 0.5%

OTHER INGREDIENTS: 99.5%

TOTAL: 100.0%

⁽¹⁾3-phenoxyphenyl methyl (4-1) isobutyl-2,4-dichloro-5-hydroxy-2,4-dimethyl-6-chloro-3-carboxylate

(IsETrans ratio: Min. 35% (4-1) cis and max. 65% (4-1) trans.)

DIRECTIONS FOR USE

For Indoor and Outdoor Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions, use restrictions and precautions.

STOCKPILE BEFORE USE

USE RESTRICTIONS

- Do not use when applying to pets or livestock, do not enter or allow others to enter until sprays have dried.
- Do not use on crops used for food, forage, or pasture.
- Do not apply this product with a sponge.
- Except when applying to pets and livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to pets and livestock, only protected handlers may be in the area during application.
- When used in Dairy Barns or Facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects.
- Remove or cover milking utensils before application. Wash teats of animals before milking.
- Do not use in outdoor residential misting systems.
- Do not use in commercial food/feed handling establishments, restaurants, or other commercial facilities where food is commercially prepared, processed, served, or stored.
- Do not use in institutions, schools, office buildings, or similar commercial buildings.
- Do not breathe vapors or spray mist.
- Cover or remove all food processing surfaces and utensils during treatment, and thoroughly wash before use.
- Do not apply when food or drinking water is present.
- Do not broadcast use on indoor residential surfaces.
- Do not use in aircraft cabins.
- Do not use in commercial animal premise automatic misting systems.
- Do not use on food or feedstuffs.
- Do not allow use indoors or outdoor drains. No permit la entrada a desagues interiores o externos.

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors. In the home, cover all food processing surfaces and seal them off, or thoroughly wash before use. Do not remove exposed food. Remove pets, birds, and other fish. Do not use on people. Do not use on people. Do not allow children or pets to contact treated areas until surfaces are dry. Do not enter or allow other to enter until sprays have dried.

Surface Spraying

Hold sprayer approximately 12 to 15 inches away from surfaces to be treated and spray until surface is damp, but not to the point of run-off.
For 60 Day Control of Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Saw-bugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Gravid Mites, Fleas, Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Dryspine Beetles, Meal Worms, Grain Mites, and Caddisflies:
Direct Fly-RID® Plus spray into hiding places, cracks and crevices, and behind shelves and drawers.

Manufactured for: **Durvet, Inc.**, 100 S.E. Magellan Drive, Blue Springs, MO 64014
Questions? Call 1-800-942-1570

To Control Silverfish:
Spray bookcases with this product.

To Control Ants:
Sprey trails, nests, and points of entry with this product.

To Control Carpet Beetles:
Sprey carpeting under carpeting and rugs. Make localized applications of this product to floor and baseboards. Spot and surface is damp, but not to the point of run-off. Spray directly into cracks, crevices and infested areas of shelving. Repeat application every 2 weeks.

To Control Bed Bugs:
Sprey mattresses lightly with this product, particularly around tufts and seams. Take beds apart and spray in all joints. Allow to dry thoroughly before remaking bed. Treat baseboards, moldings, and floors. Repeat application every 2 weeks.

To Control Fleas (Adults and Larvae) and Ticks, including Ticks which may carry Lyme Disease (Adults and Larvae):
Spot spray treatment of infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and for localized treatment of floors and floor coverings. Place fresh bedding in animal quarters following treatment. Repeat treatment every 2 weeks or 14 days as needed. Allow this product to dry thoroughly before allowing pets to contact treated areas, to control the source of flea infestations, treat your pet with a product registered for animal application. Also, treat your yard with a product registered for yard application.

To Control Clothes Moths:
Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply Fly-HP® Plus spray liberally to empty chests, closets, bureaus, and other storage areas directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping the spray nozzle at least 3 feet from fabric to avoid staining.

To Kill Lice and Loose Eggs:
Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying then proceed to spray entire area. Treaters with this product, Spray from a distance of 3 to 6 inches. Treat only the surface of the bed, and avoid spraying furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK

For Use To Protect Cattle (Beef and Dairy), Goats, Sheep, Hogs, and Horses:

To Control of Horn Flies, Face Flies, House Flies, Mosquitoes, and Gnats:

Apply a light mist of this product sufficient to wet the surface of the hair. For face flies, spray face and head. Do not spray into eyes.

To Control of Stable Flies, Horse Flies, and Deer Flies:

Apply this product at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at regular intervals to give continued protection.

To Control of Blood-Sucking Lice:

Apply this product to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To Control of Poultice Lice:

Spray roosts, walls and nests or cages thoroughly with this product. This should be followed by spraying over the birds with a fine mist.

To Control of Bed Bugs and Mites in Poultry Houses:

Spray this product into crevices of roost poles, cracks in walls, and cracks in nest where the bed bugs and mites hide.

FOR PET DOG USE: Application of product on pet dogs must only be done indoors. Consult a veterinarian before using this product on debilitated, malnourished, aged, pregnant or nursing animals. Skin irritations may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur, bathe your pet.

with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

To Control Fleas, Ticks, and Lice:

Start spraying at the ear moving the dispenser rapidly and making sure the animal's entire body is covered, including the legs and under body, with this product. While spraying, lift the hair so that the spray will penetrate into the skin. Make sure spray wets thoroughly, but does not saturate animal. Do not spray into eyes or ears. Do not spray on face or head. Use protective gloves or mitts when applying this product. Repeat every 25 days as necessary for control.

NOTE: Not for use on puppies less than 12 weeks of age.

DO NOT USE ON CATS OR KITTENS.

OUTDOOR APPLICATION

USE RESTRICTIONS

- All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 1. Application to previous surfaces such as soil, lawn, turf, and other vegetation.
 2. Perimeter band treatments of 1 foot wide or less from the base of a man-made structure to previous surfaces (e.g., soil, mulch, or lawn).
 3. Application to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure.
 4. Application around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch.
 5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level.
 6. Applications to vertical surfaces directly above previous surfaces, such as soil, lawn, turf, mulch or other vegetation only if the previous surface does not drain into ditches, storm drains, gutters, or surface waters.
- For soil or tidal applications, do not apply by ground within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply when wind speed is greater than 10 mph. For example, 2 ft by 1 ft or 4 ft by 0.5 ft.
- Do not apply when wind speed is greater than 10 mph.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during storm drains, water bodies, or aquatic habitat can occur.
- Do not spray the product into fish ponds, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during storm drains, water bodies, or aquatic habitat can occur.
- Do not allow the product to enter any drain during or after application.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot treatment or crack-and-crevice treatment.
- Do not apply or irrigate to the point of runoff.
- Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Treat surfaces to ensure thorough coverage but avoid runoff.
- To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

APPLICATION INSTRUCTIONS

For Use as a Perimeter Treatment:

To control ticks, including ticks which may carry Lyme Disease:

Use this product as a residual treatment to control ticks, including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Imnophephalus sanguineus*), American Dog Tick (*Dermacentor*

variabilis) and Western Black Legged Tick (*Ixodes pacificus*). For each of the areas listed below, apply until surface is wet, but not to the point of run-off.

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter. Apply in a band of vegetation and leaf litter 3 to 6 feet wide around the outside of buildings to help control ticks occurring there. To help a band of vegetation and leaf litter 3 to 6 feet wide around the outside of buildings to help control ticks occurring there. To help a band of vegetation and leaf litter 6 to 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

To Control Garden Insects:

For use on roses, dahlias, asters, and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites, and Leafminers. This product may be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

This product may also be used on other plants such as African violets, azaleas, begonias, camelias, carnations, chrysanthemums, delphiniums, dogwoods, English ivy, euonymus, fuchsia, geraniums, Cassia grape vines, Kentia palms, laurels, marigolds, nodotendrons, rubber plants, snapdragons, stocks, wandering jews, and zinnias.

This product is effective when used as directed against Whiteflies, Aphids, Spider Mites, Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites, and Armyworms.

When spraying plants with this product, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants, and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly with spray this product whenever possible.

Not for use on plants being grown for sale or other commercial uses, for commercial seed production, or for research purposes. Not for use in commercial greenhouses.

Resistance Management Recommendations

For resistance management, Fly-HP® Plus spray contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to Fly-HP® Plus spray and other Group 3A insecticides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Fly-HP® Plus spray or other Group 3A insecticides with different groups that control the same pests.
- Adopt an integrated pest management program for insecticide use.
- Monitor for and manage to target pest control. If the level of survival suggests the presence of resistant insects, consult with your local university's certified pest control technician.
- Contact your local extension specialist for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Cooperative Agricultural Extension Agent or your local DuPont representative at 1-800-821-5570.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry and secure area. Store in original sealed container away from food or feed.

PESTICIDE DISPOSAL: Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or other for recycling if available.

If partly filled: Call your local solid waste agency for disposal options.

DISCLAIMER

To the extent permitted by applicable law, Seller makes no representation or warranty, either express or implied, for results due to misuse, improper handling, or improper storage of this material. We does Seller assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling, or improper storage of this material.

FIRST AID

- IF ON SKIN OR ON CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for further treatment advice.

Have product container or label with you when calling a poison control center, or doctor, or going in for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-451-523-0318 (all other areas).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear the following:

- Long-sleeved shirt and long pants;
- Shoes and socks; and
- Chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or Viton.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees or other pollinating insects are foraging the treatment area.

Durvet[®], the Durvet logo and FlyRID[®] are registered trademarks of Durvet, Inc. All rights reserved.

Manufactured in the U.S.A.
with global components

ISSN:BX085
EPA Reg. No. 47000-146-17281
EPA Est. No. 47000-04-A01



PERMETHRIN GROUP 3A INSECTICIDE



Multi-Purpose Spray

Controls Flies, Mosquitoes and Gnats
35-Day Control of Fleas & Ticks on Dogs
With **CITRONELLA**

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional
Precautionary Statements and
Directions For Use.

ACTIVE INGREDIENT:	
Permethrin	0.5%
OTHER INGREDIENTS:	
TOTAL	99.5%
* <i>(3-phenoxyphenyl) methyl (+)-cis/trans-3-(2-dichloroethyl)-2,2-dimethylpyrrolidinecarboxylate</i> (cis/trans ratio: Min. 35% (+)-cis and max. 65% (+)-trans)	
100.0%	

Manufactured for: **Durvet, Inc.**, 100 S.E. Magellan Drive, Blue Springs, MD 64014
Questions? Call 1-800-821-5570

NET CONTENTS: 1 Qt. (946 mL)