

# GoatCare•2X

## TYPE C MEDICATED FEED

### Indications:

For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta* and *Trichostrongylus axei*.

### Approved For Use On:

All classes of goats.

### Active Ingredient:

Contains Rumatel® (Morantel tartrate)  
0.194% (0.88 grams per pound)

### Guaranteed Analysis:

Crude Protein (min)..... 12%  
Crude Fat (min)..... 1%  
Crude Fiber (max)..... 15%  
Acid Detergent Fiber (max)..... 23%  
Calcium (min)..... 2.8%  
Calcium (max)..... 3.3%  
Phosphorus (min)..... 0.5%  
Selenium (min)..... 0.2 PPM

### Directions for Use:

See label for full directions.

TOP-DRESS DIRECTIONS	
Body Weight (LB)	GoatCare 2X
12.5	1 oz
25	2 oz.
50	4 oz.
75	6 oz.
100	8 oz.
150	12 oz.
200	16 oz.



### Product Advantages:

- FDA Approved for use in all classes of goats
- No milk withdrawal. 30 day slaughter withdrawal.
- Convenient, single dose treatment at any time during production cycle, helping you to strategically deworm the entire herd.
- Palatable, stable in any meal form.
- Very safe. Rumatel®, when administered during safety trials at twenty times the recommended level, produced no adverse reactions. Backed by years of safe use in goats.
- Does not adversely impact reproductive performance.
- May be used frequently. Conditions of constant worm exposure may require retreatment within 2 to 4 weeks.
- 4 oz. measuring scoop included inside every bag for simple dosing.
- 3 lb bag treats 12-50 lb goats. 6 lb bag treats 24-50 lb goats.

### Packaging:

3 lb reclosable bag, 6 per case      UPC# 7-45801-10311-5  
6 lb reclosable bag, 4 per case      UPC# 7-45801-10312-2

Includes a 4 oz. measuring scoop

# GoatCare•2X

## TYPE C MEDICATED FEED

# GOAT DEWORMER

For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta* and *Trichostrongylus axei*.

#### ACTIVE INGREDIENTS:

Morantel Tartrate: 0.194% (0.88 grams per pound)

#### INGREDIENTS:

Wheat middlings, soybean hulls (max 20%), calcium carbonate, molasses products, calcium propionate (a preservative).

#### DIRECTIONS FOR USE:

- Feed as a single therapeutic treatment by feeding 0.5 pounds (0.44 grams of morantel tartrate) per 100 pounds of body weight. May be fed as the sole ration or mixed with 1 to 2 parts of complete feed or as a top dress. When used as a top dress, the medication as well as the underlying feed should be evenly distributed.
- Withhold feed overnight prior to treatment to insure the ration will be readily consumed.
- Group animals by size for optimum efficacy.
- The medicated feed should be consumed within 6 hours.
- Fresh water should be available at all times.
- After medicated feed is consumed, resume normal feeding.
- Conditions of constant worm exposure may require retreatment in 2 to 4 weeks.
- Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Under dosing may result in ineffective treatment and encourage the development of parasite resistance.

#### CAUTION:

Consult your veterinarian before using in severely debilitated animals. Do not mix in feeds containing Bentonite.

#### WARNING:

Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats. For use in goats only.

#### OTHER WARNINGS:

- Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.
- Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.
- Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of fecal egg count reduction test or another appropriate method).
- A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

#### GUARANTEED ANALYSIS:

Crude Protein (min) .....	12%
Crude Fat (min) .....	1%
Crude Fiber (max) .....	15%
Acid Detergent Fiber (max) .....	23%
Calcium (min) .....	2.8%
Calcium (max) .....	3.3%
Phosphorus (min) .....	0.5%
Selenium (min) .....	0.2 PPM

#### TOP-DRESS DIRECTIONS:

Body Weight (LB)	GoatCare•2X
12.5	1 oz
25	2 oz
50	4 oz
75	6 oz
100	8 oz
150	12 oz
200	16 oz

4 oz MEASURING SCOOP ENCLOSED.  
1 Level Scoop per 50 lb. Body Weight

ISS23XB11

Distributed by Durvet, Inc.

Blue Springs, MO 64014

www.durvet.com | 1-800-821-5570



NET WT: 3 LB (1.36 KG)



GOAT CARE-2X

Tel 800-821-5570 | Fax 816-224-3080 | info@durvet.com

010424