Canine Spectra 6

Ideal First Puppy Vaccine

6-WAY PROTECTION



SpectraTM 6 vaccine is a combination of immunogenic, attenuated strains of Canine Distemper (CDV), Canine Adenovirus Type 2 (CAV2), Canine Parainfluenza (CPIV), and Canine Parvovirus (CPV) type 2b, propagated in cell line tissue cultures. The diluent contains killed antigens for Canine Coronavirus Vaccine (CCV). The CAV2 fraction has been shown to be effective against disease due to CAV1. The CPV2b strain has been demonstrated to be effective against CPV2c in puppies with CPV maternal antibody. Gentamicin, thimerosal and amphotericin B added as preservatives.

- Comprehensive first puppy shot with Coronavirus
- Contains CPV2b strain effective protection against CPV2c
- Do-it-yourself at home
- · Easy to use vaccines complete with syringe
- How-to-vaccinate videos available online

GUARANTEED QUALITY Pure. Safe. Effective. Easy to use. Spectra & Focus vaccines are made to exacting quality standards and temperature controlled from plant to you. Peak protection against major infectious diseases. USDA approved for purity, potency, safety and efficacy.

Indications


This product has been shown to be effective for the vaccination of healthy dogs 6 weeks of age or older against canine distemper virus (CDV), infectious canine hepatitis (CAV1), canine adenovirus type 2 (CAV2), canine coronavirus (CCV), canine parainfluenza (CPIV), and canine parvovirus (CPV). The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productddata.aphis.usda.gov.

Directions and Dosage

Open the syringe by twisting or tapping the cap against a hard surface to break the heat weld. Aseptically rehydrate the vaccine cake with 1 mL of vaccine diluent supplied. SHAKE WELL. Withdraw entire contents into the syringe. Administer one 1 mL dose subcutaneously.

Subcutaneous administration: Lift the loose skin behind the neck or behind the front leg and insert needle (see illustration, arrows 1 and 2). Inject entire contents of syringe. Do not inject directly into blood vessel (see note below). IMPORTANT NOTE: Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site. All dogs should initially receive one dose of this product and a second dose

2 to 3 weeks later. The presence of maternal antibody is known to interfere with the development of active immunity in dogs and additional boosters will be required in most young animals. Historically, annual revaccination with a single dose has been recommended for this product. The need for this booster has not been established. For advice on revaccination frequency and annual booster vaccinations, consultation with a veterinarian is recommended.



Administration sites.

Precautions

Only vaccinate healthy animals. This product has not been tested in pregnant animals. Store in the dark at $35^\circ-46^\circ\text{F}$ ($2^\circ-8^\circ\text{C}$). DO NOT FREEZE. Do not mix with other products, except as specified on this label. In case of anaphylactoid reaction, administer epinephrine. Inactivate all unused contents before disposal. In case of human exposure, contact a physician.

Packaging

Single dose with syringe - 25 doses/display - 4 displays/case UPC# 7-45801-15086-7 25 doses (without syringe)/tray - 40 trays/case

25 doses (without syringe)/tray - 40 trays/ca

It's easy to do-it-yourself

Scan the code at right to view helpful how-to-vaccinate videos or visit:

VaccinateYourPets.com



