

## FOR USE IN ANIMALS ONLY

### INDICATIONS:

For use in cattle as an aid in treating uncomplicated milk fever (parturient paresis).

### DOSAGE & ADMINISTRATION:

For intravenous administration only.

**Cattle** - 250-500 mL

Dosage may be repeated at 8 to 12 hour intervals if required.

### CAUTION:

Solution should be warmed to body temperature and administered **slowly**. Entire contents should be used upon entering, as this product contains no preservative. Discard any unused portion. Do not use if solution is cloudy or contains precipitate.

### NOTE:

Monitor animal's condition closely. Re-treatment may be necessary. If there is no noticeable improvement within 24 hours following treatment, consult your veterinarian. Aseptic precautions should be observed.

NDC 30798-853-17



# Calcium Gluconate

## 23% SOLUTION

## Sterile Solution



**NET  
CONTENTS:  
500 mL  
(16.91 FL. OZ.)**

## KEEP OUT OF REACH OF CHILDREN

### ACTIVE INGREDIENTS:

**Each 500 mL Contains:**

Calcium Gluconate ..... 23% w/v  
(provides 10.71 g of Calcium\* per 500 mL)

Water for Injection ..... q.s

\*Present as salts or boryl esters of gluconic acid.

**Store between 15°C - 30°C (59°F - 86°F).**

**Bottled in the U.S.A.**

ISS24XB11

Manufactured for:

**DURVET, INC.**

Blue Springs, Missouri 64014

[www.durvet.com](http://www.durvet.com)

TAKE TIME



**OBSERVE LABEL  
DIRECTIONS**



Lot No.

Exp. Date