

## FOR ANIMAL USE ONLY

**INDICATIONS:** In cattle, horses, swine and sheep as a topical antiseptic for use prior to surgical procedures such as castrating and docking; for ringworm and foot rot; for application to the navel of newborn animals; and for minor cuts, teat sores, bruises and abrasions.

**DIRECTIONS:** Remove spray cap from 16 oz. container and refill from gallon container. Replace spray cap tightly. Hold container approximately 4 to 6 inches from the area to be sprayed. Point valve at the area to be sprayed. Pull trigger of valve and spray area once lightly. May be repeated daily, if necessary, until the area is healed.

**CAUTION:** If redness, irritation or swelling persists or increases, discontinue use and consult a veterinarian. Not for use on burns or in body cavities or deep wounds.

**NOTE:** When used on or near the teats or udders of dairy animals, the teats and udders should be thoroughly washed before the next milking to prevent contamination of milk.

Manufactured for: **DURVET, INC.**,  
Blue Springs, MO 64014 • [www.durvet.com](http://www.durvet.com)



ISS23XB07

NDC 30798-466-31



# Controlled Iodine Spray

## Antiseptic/Disinfectant



**For Skin Disinfection,  
Superficial Wounds,  
Navel Cords**

**NET CONTENTS:  
16 fl. oz. (473 mL)**

## KEEP OUT OF REACH OF CHILDREN

### INGREDIENTS:

#### Active Ingredients:

Polyethoxylated nonylphenol iodine complex  
(provides 1.00% minimum titratable iodine)

.....5.00%

**Inactive Ingredients:** .....95.00%

Purified Water and Nonionic  
Surfactant.

**CAUTION:** Harmful or fatal if  
swallowed. Do not apply to the  
eyes, mucous membranes or large  
areas of abraded skin. Avoid  
inhalation of mist.

**STORAGE:** Store at controlled  
room temperature between  
10°-30°C (50°-86°F). Keep tightly  
closed, protect from light.

Lot No.

Exp. Date

