

RECLOSABLE BAG



GoatCare•2X

TYPE C MEDICATED FEED

GOAT DEWORMER

Contains Rumatel®(Morantel Tartrate)

TREATS
12
FIFTY LB
GOATS



For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta*, and *Trichostrongylus axei*.

ACTIVE DRUG INGREDIENT
Morantel Tartrate 0.194% (0.88 grams per pound)

Rumatel is a registered trademark of Phibro Animal Health Corporation.

See Back Panel for Additional Information

NET WT: 3 LB (1.36 KG)

FOLD

FOLD



K-Seal

K-Seal

FOLD

FOLD

NET WT: 3 LB (1.36 KG)



ISS23XB11
Distributed by Durvet, Inc.
Blue Springs, MO 64014
www.durvet.com | 1-800-821-5570

Body Weight (LB)	GoatCare•2X
12.5	1 oz
25	2 oz
50	4 oz
75	6 oz
100	8 oz
150	12 oz
200	16 oz

TOP-DRESS DIRECTIONS:

Crude Protein (min)	12%
Crude Fat (min)	1%
Crude Fiber (max)	15%
Acid Detergent Fiber (max)	23%
Calcium (min)	2.8%
Calcium (max)	3.3%
Phosphorus (min)	0.5%
Selenium (min)	0.2 ppm

GUARANTEED ANALYSIS:

TREATS
12
FIFTY LB
GOATS

GOAT DEWORMER

For removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta* and *Trichostrongylus axei*.

ACTIVE INGREDIENTS:

Morantel Tartrate: 0.194% (0.88 grams per pound)

INGREDIENTS:

Wheat middlings, soybean hulls (max 20%), calcium carbonate, molasses products, calcium propionate (a preservative).

DIRECTIONS FOR USE:

• Feed as a single therapeutic treatment by feeding 0.5 pounds (0.44 grams of morantel tartrate) per 100 pounds of body weight. May be fed as the sole ration or mixed with 1 to 2 parts of complete feed or as a top dress. When used as a top dress, the medication as well as the underlying feed should be evenly distributed.

• Withdraw feed overnight prior to treatment to insure the ration will be readily consumed. • Group animals by size for optimum efficacy. • The medicated feed should be consumed within 6 hours. • Fresh water should be available at all times. • After medicated feed is consumed, resume normal feeding. • Conditions of constant worm exposure may require retreatment in 2 to 4 weeks. • Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Under dosing may result in ineffective treatment and encourage the development of parasite resistance.

CAUTION:

Consult your veterinarian before using in severely debilitated animals. Do not mix in feeds containing Bentonite.

WARNING:

Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats. For use in goats only.

OTHER WARNINGS:

• Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.

• Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

• Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of fecal egg count reduction test or another appropriate method).

• A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

RECLOSABLE BAG



GoatCare•2X
TYPE C MEDICATED FEED

↑.375" Seal Area

↑.375" Seal Area