

SEE BACK LABEL FOR DIRECTIONS FOR USE

Easy One (1) Day Treatment!

RECLOSABLE BAG

KEEP OUT OF REACH OF CHILDREN.

HORSE to FOAL 2X

Safe-Guard® (fenbendazole) 1.1%

FENBENDAZOLE TYPE B MEDICATED FEED

For the treatment and control of: **Large strongyles** (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*, *Triodontophorus* spp.), **Small strongyles** (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp.), **Pinworms** (*Oxyuris equi*), and **Ascarids** (*Parascaris equorum*).



Apple Flavored!

MULTI-DOSE

CONTENTS TREAT UP TO 2,000 LBS OF BODY WEIGHT

NET WT 1.0 LB (0.453 KG)

durvet

FOLD

↑ .25" Seal Area ↓

Mix the required amount with feed ration for easy **ONE (1) DAY** treatment!

ACTIVE DRUG INGREDIENT:

Fenbendazole: 1.1% (5 g/lb)

GUARANTEED ANALYSIS:

Crude Protein, min..... 11.50%

Crude Fat, min..... 2.50%

Crude Fiber, max..... 13.00%

INGREDIENTS:

Processed grain by-products, calcium carbonate, molasses products, roughage products, silicon dioxide, propionic acid (preservative), acetic acid, natural and artificial flavoring components (Apple Flavor), sodium propionate (preservative), potassium sorbate (preservative), propylene glycol and citric acid (preservative).

MIXING DIRECTIONS:

Mix 1 pound (contents of bag) of Type B medicated feed with 1.2 pounds of feed ingredients to manufacture Type C medicated feed containing 4540 grams (0.5%) fenbendazole per ton equivalent.

The resulting Type C feed is to be fed at the rate of 0.1 or 0.2 pounds per 100 pounds of body weight to provide 2.27 or 4.54 mg fenbendazole per pound of body weight.

For the control of large strongyles, small strongyles, and pinworms the recommended dose is 5 mg fenbendazole per kg body weight (2.27 mg fenbendazole per pound) in a **ONE (1) DAY** treatment. For the control of ascarids the recommended dose is 10 mg fenbendazole per kg body weight (4.54 mg fenbendazole per pound) in a **ONE (1) DAY** treatment.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Do not use in horses intended for human consumption.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

- **Apple flavored pellets**
- **For Horses of All Ages**

Learn more about **Horse to Feed ZX**
and Durvet's line of apple flavored
equine dewormers at:

www.Durvet.com

Approved by FDA under NADA # 131-675

Safe-Guard® is a registered trademark of Intervet International B.V.

durvet
www.durvet.com

Visit us online!



Like us on



facebook.com/Durvet

Manufactured for: Durvet, Inc.
Blue Springs, MO 64014
1-800-821-5570
www.durvet.com

ISS23XB08



NET WT 1.0 LB (0.453 KG)