

durvet

IVERMECTIN POUR-ON
(ivermectin topical solution)



for Cattle

Contains 5 mg ivermectin / mL

Parasiticide

Kills:

Roundworms, (including Brown Stomach Worm), Lungworms, Grubs,
Sucking Lice, Biting Lice, Mange Mites, Horn Flies.

Manufactured for:

DURVET, INC.

Blue Springs, Missouri 64014

www.durvet.com

Approved by FDA under ANADA # 200-272



7 45801 11045 8

169 fl oz (5 L)

Rev. 03/20
010215L03

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS:

For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle. See package insert for complete indications and use directions.

Recommended Dose: 1 mL per 22 lb of body weight.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

WARNING

NOT FOR USE IN HUMANS.

Keep this and all drugs out of the reach of children.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Norbrook at 1-866-591-5777. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

WARNING! FLAMMABLE!
KEEP AWAY FROM HEAT,
SPARKS, OPEN FLAME, AND
OTHER SOURCES OF IGNITION.

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin.

To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

RESIDUE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

PRECAUTIONS:

Store at or below 25°C (77°F), excursions permitted up to 40°C (104°F). Protect from light.

Use only in well-ventilated areas or outdoors. Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced.

Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when Ivermectin Pour-On (ivermectin topical solution) is stored at temperatures below 32°F. Allowing it to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g. caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermectin Pour-On is not recommended for use in species other than cattle.

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Restricted Drug - California. Use Only as Directed.

Approved by FDA under ANADA # 200-272

Made in the UK.

Manufactured by:
DURVET, INC.
Blue Springs, Missouri 64014
www.durvet.com

169 fl oz (5 L)

Lot No. & Exp. Date ▼ 01215104 Rev. 03/20