



Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS:
For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice, and sarcoptic mange mites in cattle.

See package insert for complete indications and use directions.

DOSAGE:
The dose rate is 1 mL for each 22 lb of bodyweight. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

ADMINISTRATION:
Connect the applicator gun to the collapsible pack as follows:
Attach the open end of the draw-off tubing to the dosing equipment. (Because of the solvents used in the formulation, only the applicator gun from Simcro Tech, or equivalent, is recommended. Other applicators may exhibit compatibility problems, resulting in locking, incorrect dosage or leakage.)
Replace the shipping cap with the draw-off cap and tighten down. Attach draw-off tubing to the draw-off cap.
Gently prime the applicator gun, checking for leaks.
Follow the manufacturer's direction for adjusting the dose.
When the interval between uses of the applicator gun is expected to exceed 12 hours, disconnect the gun and draw-off tubing from the product container and empty the product from the gun and tubing back into the product container. To prevent removal of special lubricants from the Simcro Tech gun, the gun and tubing must not be washed.

Manufactured for:
DURVET, INC.
Blue Springs, MO 64014
www.durvet.com

Made in the UK.



ANIMAL SAFETY:
Studies conducted in the U.S.A. have demonstrated the safety margin for ivermectin. Based on plasma levels, the topically applied formulation is expected to be at least as well tolerated by breeding animals as is the subcutaneous formulation which had no effect on breeding performance.

WARNING
NOT FOR USE IN HUMANS
Keep this and all drugs out of the reach of children.

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

WARNING! FLAMMABLE!
KEEP AWAY FROM HEAT, SPARKS,
OPEN FLAME, AND OTHER SOURCES
OF IGNITION.

RESIDUE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

OTHER WARNINGS
Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.
Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method).

A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.



Contains 5 mg ivermectin/mL

169 fl oz (5 L)



Contains 5 mg ivermectin / mL

Parasiticide

Kills: Roundworms (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies.

Contains 200 - 550 lb doses

Approved by FDA under ANADA # 200-272

169 fl oz (5 L)



This topically applied formulation of Ivermectin Pour-On (ivermectin topical solution) delivers internal and external parasite control in one convenient low-volume application.

PRECAUTIONS:
Store at or below 25°C (77°F), excursions permitted up to 40°C (104°F). Protect from light.

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced.

Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when Ivermectin Pour-On is stored at temperatures below 32°F. Allowing it to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g. caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermectin Pour-On is not recommended for use in species other than cattle.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Norbrook at 1-866-591-5777. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

Restricted Drug - California. Use Only as Directed.

Rev. 03/20
6215869C06