For Intramuscular Use in Swine Over 300 lbs.
Restricted Drug - Use Only As Directed (California). For Use in Animals Only.

Dosage: Usual daily dose for arthritis or mycoplasma pneumonia - 5 mg per pound of body weight (1 mL per each 60 pounds of body weight) intramuscularly for three to seven days. See package insert for complete product information.

Contains per mL: Lincomycin Hydrochloride equivalent to Lincomycin, 300 mg; also Benzyl Alcohol 9.45 mg added as preservative.

Store at controlled room temperature: 20° to 25°C (68° to 77°F) [see USP].

L-4636-04 Rev. 12-16

Lot No. Exp. Date

7 45801 10816 5

NDC 30798-336-10

durvet

LINCOMYCIN 300

300 mg/mL
Sterile Lincomycin Hydrochloride Injection

Swine Antibiotic

NET CONTENTS:
100 mL (3.3 fl oz)

Indicated for the treatment of arthritis caused by susceptible organisms and for mycoplasma pneumonia.

Warning: Not for human use. Keep out of reach of children. Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

TAKE TIME OBSERVE LABEL DIRECTIONS

ANADA#: 200-315 Approved by F.D.A.

Manufactured for
DURVET, INC
Blue Springs, Missouri 64014
www.durvet.com
ISS16XB12

Manufactured by
Sparhawk Laboratories, Inc.
Lenexa, KS 66215, USA

OPEN HERE

9.45 mg added as preservative. Supplied in 100 mL vials.

25 mg/mL: Special baby pig concentration. Each mL contains lincomycin hydrochloride equivalent to lincomycin, 25 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

Indicated for the treatment of arthritis caused by susceptible organisms and for mycoplasma pneumonia.

Warning: Not for human use. Keep out of reach of children. Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

TAKE TIME OBSERVE LABEL DIRECTIONS

ANADA#: 200-315 Approved by F.D.A.

Manufactured for
DURVET, INC
Blue Springs, Missouri 64014
www.durvet.com
ISS16XB12

Manufactured by
Sparhawk Laboratories, Inc.
Lenexa, KS 66215, USA

Lincomycin Injectable

brand of Lincomycin hydrochloride injectable

For Intramuscular Use in Swine Only

Lincomycin Injectable contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis var. lincolnensis*, which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

INDICATIONS FOR SWINE

Lincomycin Injectable is indicated for the treatment of infectious forms of arthritis

caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp. It is also indicated for the treatment of mycoplasma pneumonia.

CONTRAINDICATIONS

As with all drugs, the use of Lincomycin Injectable is contraindicated in animals previously found to be hypersensitive to the drug.

WARNING

Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment. **Not for human use.**

CAUTION

If no improvement is noted within 48 hours, consult a veterinarian.

ADVERSE REACTIONS

The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert

to the possibility that it may occur. Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION

For arthritis or mycoplasma pneumonia - 5mg per pound of body weight intramuscularly once daily for three to seven days as needed.

When using Lincomycin Injectable containing 25 mg/mL, 1 mL/5 lb. body weight will provide 5 mg/lb.

When using Lincomycin Injectable

containing 100 mg/mL, 1 mL/20 lb. body weight will provide 5 mg/lb.

When using Lincomycin Injectable containing 300 mg/mL, 1 mL/60 lb. body weight will provide 5 mg/lb.

For optimal results, initiate treatment as soon as possible. As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

HOW SUPPLIED

Lincomycin Injectable is available in three concentrations: 300 mg/mL, 100 mg/mL, and 25 mg/mL.

300 mg/mL: For use in swine weighing 300 pounds or more. Each mL contains lincomycin hydrochloride equivalent to lincomycin 300 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

100 mg/mL: Each mL contains lincomycin hydrochloride equivalent to lincomycin 100 mg; also Benzyl Alcohol,