

FOR ANIMAL USE ONLY

INDICATIONS: For use as an aid in prevention and treatment of Ketosis (Acetonemia) in dairy cattle.

DIRECTIONS FOR USE: For use mixed with feed or as a drench.

Prevention: Dairy Cattle: 6-8 fl oz/head/day
Begin 2 weeks before calving and continue for 6 weeks after calving.

Treatment: Dairy Cattle: 6-16 fl oz/head/day
Dosage dependant on severity, to be given for 10 days.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Manufactured for: **DURVET, INC.**,
Blue Springs, MO 64014 • www.durvet.com ISS23XB08

NDC 30798-464-35



Propylene Glycol

For Use as an Aid in Prevention
and Treatment of Ketosis



**NET CONTENTS:
1 Gallon (3.785 L)**

KEEP OUT OF REACH OF CHILDREN

Consult veterinarian if condition does not respond.

LIVESTOCK DRUG

STORAGE: Store at controlled room temperature between 2°-30°C (36°-86°F). Keep container tightly closed, when not in use.

Seller makes no warranty of any kind whether in accordance with directions or not.

ACTIVE INGREDIENTS:

Propylene Glycol100%

Lot No.

Exp. Date

