

ADMINISTRATION:
After removing the cap, the paste should be deposited on the dorsum of the tongue. Introduce the nozzle end of the syringe at the corner of the mouth. Direct the syringe backwards and depress the plunger to deposit the paste onto the tongue. Given in this manner, it is unlikely that rejection of the paste will occur. Raising the horse's head sometimes assists in the swallowing process. When only part of the paste has been used, replace the cap on the syringe nozzle.

EFFICACY:
Critical (worm count) studies in horses demonstrated that pyrantel pamoate paste administered at the recommended dosage was efficacious against mature infections of *Strongylus vulgaris* (>90%), *S. edentatus* (69%), *S. equinus* (>90%), *Oxyuris equi* (81%), *Parascaris equorum* (>90%), and small strongyles (>90%).

WARNINGS
For oral animal use only. Do not use in horses intended for human consumption.

HUMAN WARNINGS:
Keep out of the reach of children.

It is recommended that severely debilitated animals not be treated with this preparation.

OTHER WARNINGS: Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's

effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

RECOMMENDED STORAGE: Store at 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 30°C (59°F - 86°F).



Approved by FDA under ANADA # 200-350

PYRANTEL PASTE
(pyrantel pamoate)
HORSE DEWORMER

NDC 96798-836-83



Net Contents:
One syringe containing 3.6 grams of pyrantel base in 23.6 grams of paste.

Apple Flavored



& Copyrighted by

PYRANTEL PASTE
(pyrantel pamoate)

Apple Flavored



Net Weight: 23.6 g (0.83 oz).
Each mL contains 171 mg pyrantel base as pyrantel pamoate.



PYRANTEL PASTE
(pyrantel pamoate)



Manufactured by: Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2M4
Distributed by: durvet Inc.
Blue Springs, MD 64014
www.durvet.com
Rev. 01/21 18PYR016D

Body Weight Range	Volume	mg Pyrantel Base
up to 300 lb	1/4 syringe	900 mg
301 - 600 lb	1/2 syringe	1800 mg
601 - 900 lb	3/4 syringe	2700 mg
901 - 1200 lb	1 full syringe	3600 mg

DOSE: Position ring-gauge over appropriate mark on plunger. Each mL contains 171 mg of pyrantel base as pyrantel pamoate.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

For maximum control of parasites, it is recommended that foals (2-8 months of age) be dosed every 4 weeks for a minimum of the potential source of infection that the mare may pose to the foal, the mare should be treated 1 month prior to anticipated foaling date followed by retreatment 10 days to 2 weeks after birth of foal. Horses and ponies over 8 months of age should be routinely dosed every 6 weeks.

Chemical name: Ethyl 5-benzothiazol-2-yl-1-methyl-2-(2-thienyl)-vinyl-pyridine-4,4'-methylenebis[3-hydroxy-2-naphtholate] (1:1)

DESCRIPTION: Pyrantel Paste (pyrantel pamoate) is a pale yellow to buff paste containing 43.9% w/w pyrantel pamoate in an inert vehicle. Each syringe contains 3.6 grams of pyrantel base in 23.6 grams paste. Each mL contains 171 mg pyrantel base as pyrantel pamoate.

COMPOSITION: Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The pyrantel base per lb of body weight. The syringe has 4 weight mark increments. Pyrantel base (pyrantel pamoate) is to be administered as a single oral dose of 3 mg of pyrantel base per lb of body weight.

INDICATIONS FOR USE: For the removal and control of mature infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*, small strongyles, pinworms (*Oxyuris equi*, and large roundworms (*Parascaris equorum*) in horses and ponies. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSE AND TREATMENT: Pyrantel Paste (pyrantel pamoate) is to be administered as a single oral dose of 3 mg of pyrantel base per lb of body weight. The syringe has 4 weight mark increments. Each weight mark indicates the recommended dose for 300 lb of body weight.

Read Entire Carton Carefully Before Using This Product

PYRANTEL PASTE
(pyrantel pamoate)