FOR USE IN ANIMALS ONLY

ADMINISTRATION AND DOSAGE:

Intramuscular or subcutaneous administration only. If dose is greater than 5 mL, equally divide the dose and inject at two different sites.

SUGGESTER	DOCACE.	

CALLE (Dally allu bee	1)
Cows and Heifers	
2-3 wks pre-part	tum
At calving	8-10

End of lactation	8-10
Calves	
At birth	4-6
At weaning	4-6
Yearlings	5-6

CAUTION: Do not exceed recommended dosage. Occasionally, reactions of an anaphylactic or allergic nature may occur. Should such reactions occur, treat immediately with injection of epinephrine or antihistamines. Administration of this product to well nourished animals may cause hypervitaminosis D which may result in hypercalcemia.

WARNING: Do not add water to solution.

Manufactured for: **DURVET, INC.** Blue Springs, Missouri 64014 www.durvet.com



Vitamin E+AD

INJECTABLE TOCOPHEROL + AD (Vitamin A, D₃, and E)

INTRAMUSCULAR OR SUBCUTANEOUS INJECTION ONLY



KEEP OUT OF REACH OF CHILDREN

Each mL contains:

ACTIVE INGREDIENTS: 300 IU's of Vitamin E (as d-alphatocopherol, a natural source of Vitamin E); 100,000 IU's Vitamin A Propionate; 10,000 IU's Vitamin D₃,

INACTIVE INGREDIENTS: 2% Benzyl Alcohol (preservative) in a water emulsifiable base.

INDICATIONS: Vitamin E+AD is a clear, sterile, non-aqueous solution of Vitamin A, Vitamin Ds, and Vitamin E. The product is intended as a supplemental source of vitamin A, D, and E in cattle.

Natural Vitamin A (carotenes), and tocopherols in feedstuffs can be destroyed through processing, ensiling and storage. Due to these losses, reduced intakes of fat-soluble vitamins can occur in animals amintained in continual confinement compared to animals allowed to graze lush pasture, Intramuscular or subcutaneous injections offer an efficient and rapid method to increase vitamin A, vitamin D and vitamin E status of animals.

Store between 15° and 30°C (59°-86°F) in a dark place.

V-3234-05 REV 08-16

Lot No.: EXP. DATE:

