

## FOR USE IN ANIMALS ONLY

### ADMINISTRATION AND DOSAGE:

Intramuscular or subcutaneous administration only. If dose is greater than 5 mL, equally divide the dose and inject at two different sites.

### SUGGESTED DOSAGE:

#### SWINE

	mL	International Units (I.U.)
Sows and Gilts		
2 wks pre-partum	4-6	1200-1800
2 wks pre-breeding	4-6	1200-1800

#### Pigs

At birth	1-2	300-600
At weaning	2-3	600-900

#### CATTLE (Dairy and Beef)

Cows and Heifers		
2-3 wks pre-partum	8-10	2400-3000
At calving	8-10	2400-3000

#### Calves

At birth	4-6	1200-1800
At weaning	4-6	1200-1800

#### Yearlings

	5-6	1500-1800
--	-----	-----------

#### SHEEP

##### Ewes

2-3 wks pre-partum	4-5	1200-1500
At lambing	4-5	1200-1500

##### Lambs

At birth	2-3	600-900
At weaning	2-3	600-900

##### Finishing Lambs

	3-4	900-1200
--	-----	----------

Manufactured for:

**DURVET, INC.**

Blue Springs, Missouri 64014

[www.durvet.com](http://www.durvet.com)

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

V-4363-05

NDC 30798-694-13

**durvet**

# Vitamin E 300

**INJECTABLE TOCOPHEROL**  
( 300 I.U. Vitamin E per mL )

**INTRAMUSCULAR OR  
SUBCUTANEOUS INJECTION ONLY**



**NET CONTENTS:**  
**250 mL**  
**(8.5 FL. OZ.)**

## KEEP OUT OF REACH OF CHILDREN

**Each mL contains:**

**ACTIVE INGREDIENTS:** 300 IU's of Vitamin E (as d-alpha-tocopherol, a natural source of Vitamin E).

**INACTIVE INGREDIENTS:** 2% Benzyl Alcohol (preservative) in a water emulsifiable base.

**INDICATIONS:** Injectable Tocopherol (natural-source vitamin E) is a clear, sterile, non-aqueous solution of d-alpha-tocopherol for use as an aid in the prevention and treatment of vitamin E deficiencies in swine, cattle, and sheep.

The product is intended as a supplemental source of natural vitamin E.

Natural tocopherols in feedstuffs can be destroyed through processing, ensiling and storage. A reduced vitamin E intake can result in marginal deficiencies that may not be visible.

Intramuscular or subcutaneous injections offer an efficient and rapid method to increase vitamin E status of animals.

**WARNING:** Do not add water to solution. Not for human use.

Keep out of reach of children. **Store between 15° and 30°C (59°-86°F) in a dark place.**

**CAUTION:** Do not exceed recommended dosage. Occasionally, reactions of an anaphylactic or allergic nature may occur. Should such reactions occur, treat immediately with injection of epinephrine or antihistamines.

REV 01-17

Lot No.:

EXP. DATE:

